

Requisition #: 9900001
Patient Name: Report Sample
Date of Birth: 03/09/1960 **Patient Age:** 65
Patient Sex: F

Practitioner: NO PHYSICIAN
Date of Collection: 04/05/2025
Time of Collection: 10:00 AM
Report Date: 06/19/2025












Microbial Organic Acids Test






Metabolic Markers in Urine Reference Range (mmol/mol creatinine) Patient Value Reference Population - Females Age 13 and Over

Intestinal Microbial Overgrowth





Yeast and Fungal Markers

1 Citramalic	≤ 3.6	H	12	
2 5-Hydroxymethyl-2-furoic	≤ 14		2.0	
3 3-Oxoglutaric	≤ 0.33	H	4.0	
4 Furan-2,5-dicarboxylic	≤ 16	H	63	
5 Furancarbonylglycine	≤ 1.9	H	6.0	
6 Tartaric	≤ 4.5	H	7.0	
7 Arabinose	≤ 29		8.0	
8 Carboxycitric	≤ 29		12	
9 Tricarballic	≤ 0.44	H	1.0	



Bacterial Markers

10 Hippuric	≤ 613		9.0	
11 2-Hydroxyphenylacetic	0.06 - 0.66	H	3.0	
12 4-Hydroxybenzoic	≤ 1.3		1.0	
13 4-Hydroxyhippuric	0.79 - 17		3.0	
14 DHPPA (Beneficial Bacteria)	≤ 0.38	H	2.0	

Clostridia Bacterial Markers

15 4-Hydroxyphenylacetic (<i>C. difficile</i> , <i>C. stricklandii</i> , <i>C. lituseburens</i> & others)	≤ 19		5.0	
16 HPPHA (<i>C. sporogenes</i> , <i>C. caloritolerans</i> , <i>C. botulinum</i> & others)	≤ 208		10	
17 4-Cresol (<i>C. difficile</i>)	≤ 75		6.0	
18 3-Indoleacetic (<i>C. stricklandii</i> , <i>C. lituseburens</i> , <i>C. subterminale</i> & others)	≤ 11		11	

Additional Indicators

19 3-Hydroxy-3-methylglutaric	0.17 - 39	L	0	
20 2-Hydroxyhippuric	≤ 1.3		1.0	

This test was developed, and its performance characteristics determined by Mosaic Diagnostics Laboratory. It has not been cleared or approved by the US Food and Drug Administration, however, does comply with CLIA regulations for clinical use.

The results should be interpreted in conjunction with the complete clinical picture, given patient history and presentation, and at the discretion of the medical provider.

Methodology: GC-MS. Creatinine by Jaffe Reaction. *The creatinine test is performed to adjust metabolic marker results for differences in fluid intake. Urinary creatinine, from a random collection, has limited diagnostic value due to variability as a result of recent fluid intake.

Microbial Organic Acids Test



6-2025 Rev 2 Page 1 of 3

Requisition #: 9900001
Patient Name: Report Sample

Practitioner: NO PHYSICIAN
Date of Collection: 04/05/2025

Indicator of Fluid Intake

21 *Creatinine 100 mg/dL

*The creatinine test is performed to adjust metabolic marker results for differences in fluid intake. Urinary creatinine has limited diagnostic value due to variability as a result of recent fluid intake. Samples are rejected if creatinine is below 20 mg/dL unless the client requests results knowing of our rejection criteria.

Explanation of Report Format



The reference ranges for organic acids were established using samples collected from typical individuals of all ages with no known physiological or psychological disorders. The ranges were determined by calculating the mean and standard deviation (SD) and are defined as $\pm 2SD$ of the mean. Reference ranges are age and gender specific, consisting of Male Adult (≥ 13 years), Female Adult (≥ 13 years), Male Child (<13 years), and Female Child (<13 years).

There are two types of graphical representations of patient values found in the new report format of both the standard Organic Acids Test and the Microbial Organic Acids Test.

The first graph will occur when the value of the patient is within the reference (normal) range, defined as the mean plus or minus two standard deviations.

The second graph will occur when the value of the patient exceeds the upper limit of normal. In such cases, the graphical reference range is "shrunk" so that the degree of abnormality can be appreciated at a glance. In this case, the lower limits of normal are not shown, only the upper limit of normal is shown.

In both cases, the value of the patient is given to the left of the graph and is repeated on the graph inside a diamond. If the value is within the normal range, the diamond will be outlined in black. If the value is high or low, the diamond will be outlined in red.

Example of Value Within Reference Range**Example of Elevated Value**

Requisition #: 9900001
Patient Name: Report Sample

Practitioner: NO PHYSICIAN
Date of Collection: 04/05/2025

Interpretation

High yeast/fungal metabolites (Markers 1,2,3,4,5,6,7,8) indicate a yeast/fungal overgrowth of the gastrointestinal tract. Prescription or natural (botanical) anti-fungals, along with supplementation of high potency multi-strain probiotics (20-50 billion cfu's), may reduce yeast/fungal levels.

High tricarballic acid (propane-1,2,3-tricarboxylic acid) (Marker 9) could be caused by the intake of corn or corn-based food contaminated with fumonisins, a group of mycotoxins produced primarily by *F. verticillioides*, and other related species. Tricarballic acid is released from fumonisins during passage through the gastrointestinal tract. Tricarballic acid is an inhibitor of the enzyme aconitase and therefore interferes with the Krebs cycle. The main symptoms of aconitase deficiency are myopathy and exercise intolerance. It may also act as a magnesium chelator. Tricarballic acid is also metabolite of a component of a substance in modified corn starch, octenylsuccinic acid, found in a number of infant formulas such as Nutramigen, Vivonex, and Pregestimil. In addition, tricarballic acid is a byproduct of beet sugar and maple sugar refining and might appear after ingestion of these sugars. Tricarballic acid is also released from fumonisins upon certain food processing conditions. Clinical syndromes due to the intact mycotoxin are rare and characterized by abdominal pain and diarrhea. A specific role for fumonisins in the development of neural tube defects was suggested after the appearance of a cluster of such defects in Texas associated with consumption of corn from the heavily fumonisin-contaminated 1989 corn crop. More recent studies have shown that fumonisin B1 inhibits folate metabolism in cultured cells.

High 2-hydroxyphenylacetic acid (Marker 11) is associated with intestinal bacteria overgrowth and with the genetic disease phenylketonuria (PKU).

High DHPPA (3,4 dihydroxyphenylpropionic acid) (Marker 14) indicates excessive intake of chlorogenic acid, a common substance found in beverages and in many fruits and vegetables, including apples, pears, tea, coffee, sunflower seeds, carrots, blueberries, cherries, potatoes, tomatoes, eggplant, sweet potatoes, and peaches. Harmless or beneficial bacteria such as *Lactobacilli*, *Bifidobacteria*, and *E. coli* mediate the breakdown of chlorogenic acid to 3,4-dihydroxyphenylpropionic acid (DHPPA), and high values may indicate increased amounts of these species in the GI tract. In addition, one *Clostridia* species, *C. orbiscindens*, can convert the flavanoids luteolin and eriodictyol, occurring only in a relatively small food group that includes parsley, thyme, celery, and sweet red pepper to 3,4-dihydroxyphenylpropionic acid. The quantity of *Clostridia orbiscindens* in the GI tract is negligible (approximately 0.1% of the total bacteria) compared to the predominant flora of *Lactobacilli*, *Bifidobacteria*, and *E. coli*. Consequently, this marker is essentially useless as a general *Clostridia* marker but may be a good indicator of the presence of beneficial flora.

High quality nutritional supplements can be purchased through your practitioner or at New Beginnings Nutritionals, www.NBNUS.com <<http://www.NBNUS.com>>, or call 877-575-2467.

The nutritional recommendations in this test are not approved by the US FDA. Supplement recommendations are not intended to treat, cure, or prevent any disease and do not take the place of medical advice or treatment from a healthcare professional.